ELSEVIER

Environmental Pollution 149 (2007) I–IV

ENVIRONMENTAL POLLUTION

ww.elsevier.com/locate/envpol

Contents of Volume 149

Number 1 2007

Regular Papers

1 **Modelling phosphate adsorption to the soil: Application of the non-ideal competitive adsorption model**
J.S. Abou Nohra, C.A. Madramootoo, W.H. Hendershot

The NICA model accurately described the adsorption of phosphate to some southern Quebec soils.

10 **Metal contamination of vineyard soils in wet subtropics (southern Brazil)**
N. Mirlean, A. Roisenberg, J.O. Chies

Copper-based pesticide use in wet subtropics is environmentally more risky.

18 **Impaired terrestrial and arboreal locomotor performance in the western fence lizard (*Sceloporus occidentalis*) after exposure to an AChE-inhibiting pesticide**
S.E. DuRant, W.A. Hopkins, L.G. Talent

Exposure to an acetylcholinesterase-inhibiting pesticide alters locomotor performance in western fence lizards.

25 **Polychlorinated biphenyls, organochlorinated pesticides, and polybrominated diphenyl ethers in the cerebral cortex of wild river otters (*Lontra canadensis*)**
N. Basu, A.M. Scheuhammer, M. O'Brien

River otters accumulated PCBs, OCPs, and PBDEs, but at levels below thresholds for neurotoxic effects.

31 **Fluxes of oxidised and reduced nitrogen above a mixed coniferous forest exposed to various nitrogen emission sources**
J. Neirynck, A.S. Kowalski, A. Carrara, G. Genouw, P. Berghmans, R. Ceulemans

Reduced nitrogen was found to be the main contributor to total deposition which was predominantly governed by dry deposition.

44 **Effects of metals on life cycle parameters of the earthworm *Eisenia fetida* exposed to field-contaminated, metal-polluted soils**
J. Nahmani, M.E. Hodson, S. Black

Soil pH, organic carbon content and texture can exert a greater influence on earthworm life cycle parameters than soil metal concentrations at metal-contaminated sites.

59 **Fate of metal-associated POM in a soil under arable land use contaminated by metallurgical fallout in northern France**
J. Labanowski, J. Sebastia, E. Foy, T. Jongmans, I. Lamy, F. van Oort

Small POM-derived organic particles may favor long-term metal sequestration in soils.

70 **Effect of organic suspended solids and their sedimentation on the surrounding sea area**
M. Fukue, C.N. Mulligan, Y. Sato, T. Fujikawa

Suspended solids can act as adsorbents for hazardous substances in surface water and their sedimentation can prevent the pollutants from spreading into the ocean.

79 **Fate and effects of heavy metals in salt marsh sediments**
K. Suntornvongsagul, D.J. Burke, E.P. Hamerlynck, D. Hahn

Spartina patens accumulates heavy metals in roots without significant translocation into shoots, and with only small seasonal effects on plant growth performance and mycorrhizal colonization.

92 **Metal accumulation in the polychaete *Hediste japonica* with emphasis on interaction between heavy metals and petroleum hydrocarbons**
F.-H. Sun, Q.-X. Zhou

The influences of petroleum hydrocarbons on Cd and Cu accumulation in H. japonica depend on their concentration combinations and exposure time.

99 **Alfalfa nutritive quality for ruminant livestock as influenced by ambient air quality in west-central Alberta**
J.C. Lin, M. Nosal, R.B. Muntifering, S.V. Krupa

Exposure to ambient concentrations of phytotoxic air pollutants affected nutritive quality of alfalfa for ruminant livestock in a yield-dependent manner.

104 **Distribution of polycyclic aromatic hydrocarbons (PAHs) and tributyltin (TBT) in Barcelona harbour sediments and their impact on benthic communities**
X. Martínez-Lladó, O. Gibert, V. Martí, S. Díez, J. Romo, J.M. Bayona, J. de Pablo

A possible correlation exists between TBT concentration in sediments and ecological effects on benthos in Barcelona harbour.

doi:10.1016/S0269-7491(07) 00423-X

114 Butyltins, trace metals and morphological variables in surf scoter (*Melanitta perspicillata*) wintering on the south coast of British Columbia, Canada
J.E. Elliott, M.L. Harris, L.K. Wilson, B.D. Smith, S.P. Batchelor, J. Maguire

Hepatic butyltin concentrations in surf scoters increased significantly over a winter in the Strait of Georgia, and correlated negatively with body condition.

Short communication

125 Synthesis, characterization and application of a novel chemical sand-fixing agent-poly(aspartic acid) and its composites
J. Yang, F. Wang, L. Fang, T. Tan

A novel biodegradability polymer significantly improved sand particles' compressive strength and anti-wind erosion properties.

Number 2 2007

Mini-Review

131 Export of toxic chemicals – A review of the case of uncontrolled electronic-waste recycling
M.H. Wong, S.C. Wu, W.J. Deng, X.Z. Yu, Q. Luo, A.O.W. Leung, C.S.C. Wong, W.J. Luksemburg, A.S. Wong

Uncontrolled e-waste recycling is ruining the environment.

Regular Papers

141 Declining metal levels at Foundry Cove (Hudson River, New York): Response to localized dredging of contaminated sediments
J.A. Mackie, S.M. Natali, J.S. Levinton, S.A. Sañudo-Wilhelmy

Dredging of a hotspot of metal-contaminated sediment is associated with a recognizable local and river-wide decline in cadmium in the Hudson River, New York.

149 Control of lead solubility in soil contaminated with lead shot: Effect of soil pH
C.P. Rooney, R.G. McLaren, L.M. Condron

Sorption processes control Pb^{2+} ion activity in soils contaminated with Pb shot.

158 Heavy metals in coastal wetland sediments of the Pearl River Estuary, China
Q.S. Li, Z.F. Wu, B. Chu, N. Zhang, S.S. Cai, J.H. Fang

We found that the sediment in coastal wetlands of the Pearl River Estuary was polluted by Cd, Zn and no longer suitable for the current wetland utilization strategies.

165 Effect of amendment C:N ratio on plant richness, cover and metal content for acidic Pb and Zn mine tailings in Leadville, Colorado
S. Brown, P. DeVolder, H. Compton, C. Henry

Increasing the carbon to nitrogen ratio of a soil amendment to >20:1 increased plant species richness for high metal mine tailings.

173 Tree species effect on the redistribution of soil metals
J. Mertens, L. Van Nevel, A. De Schrijver, F. Piesschaert, A. Oosterbaan, F.M.G. Tack, K. Verheyen

Trees (33-year-old) growing on polluted dredged sediment have influenced the metal concentration in the upper soil layer and there was a significant tree species effect.

182 Bioaccumulation of trace elements in omnivorous amphibian larvae: Implications for amphibian health and contaminant transport
J.M. Unrine, W.A. Hopkins, C.S. Romanek, B.P. Jackson

Omnivorous amphibian larvae can be efficient accumulators of trace elements.

193 Remobilization of polycyclic aromatic hydrocarbons during the resuspension of Yangtze River sediments using a particle entrainment simulator
J. Feng, Z. Yang, J. Niu, Z. Shen

Researches on remobilization of PAHs during sediment resuspension provide insight into how sediment bound PAHs release to the water column.

201 Fluxes of inorganic and organic arsenic species in a Norway spruce forest floor
J.-H. Huang, E. Matzner

The forest floor layers are generally a source for inorganic arsenic species but a sink for most organic arsenic species under the present deposition rate.

209 Responses of the mussel *Anodontites trapesialis* (Unionidae) to environmental stressors: Effect of pH, temperature and metals on filtration rate
R. Loayza-Muro, R. Elías-Letts

The freshwater mussel Anodontites trapesialis was used as a novel ecotoxicological test organism for the Amazonian region of Peru.

216 The impact of episodic coal mine drainage pollution on benthic macroinvertebrates in streams in the Anthracite region of Pennsylvania
A. MacCausland, M.E. McTammany

Episodic coal mine pollution decreases benthic macroinvertebrate richness and density.

227 Effects of chemical additives on hydrocarbon disappearance and biodegradation in freshwater marsh microcosms
J.A. Nyman, P.L. Klerks, S. Bhattacharyya

Hydrocarbon disappearance and biodegradation were insensitive to common commercial additives.

239 **Active biomonitoring of heavy metal pollution using *Rosa rugosa* plants**
G.L. Calzoni, F. Antognoni, E. Pari, P. Fonti, A. Gnes, A. Speranza

Rosa rugosa leaves and pollen are – to different extents – suitable monitoring tools for heavy metal pollution.

Short communication

246 **The impact of sewage irrigation on the uptake of mercury in corn plants (*Zea mays*) from suburban Beijing**
S.E. Rothenberg, X. Du, Y.-G. Zhu, J.A. Jay

Hg content in the roots of corn plants (Zea mays) was positively correlated with soil Hg content, but negatively correlated with Hg content in stems.

Erratum

252 **Erratum to "Diclofenac disposition in Indian cow and goat with reference to *Gyps* vulture population declines" [Environmental Pollution 147 (2007) 60–65]**
M.A. Taggart, R. Cuthbert, D. Das, C. Sashikumar, D.J. Pain, R.E. Green, Y. Feltrer, S. Shultz, A.A. Cunningham, A.A. Meharg

Number 3 2007

SPECIAL ISSUE: Air Pollution and Vegetation Effects Research in National Parks and Natural Areas: Implications for Science, Policy and Management

Commentary

253 **Commentary for papers resulting from the recent symposium on air pollution and vegetation effects in national parks and natural areas: Implications for science, policy and management**
H.S. Neufeld, A.H. Chappelka

256 **Air quality in natural areas: Interface between the public, science and regulation**
K.E. Percy, D.F. Karnosky

Appropriate endpoints and exposure–response indices can improve assessment of air pollutant risk to forests in natural areas.

268 **Translating science into policy: Using ecosystem thresholds to protect resources in Rocky Mountain National Park**
E. Porter, S. Johnson

A collaborative approach between scientists and policymakers is described for addressing nitrogen deposition effects to Rocky Mountain National Park, USA.

281 **Estimates of critical acid loads and exceedances for forest soils across the conterminous United States**
S.G. McNulty, E.C. Cohen, J.A. Moore Myers, T.J. Sullivan, H. Li

This simple mass balance equation estimated that 17% of US forest soils exceed their critical acid load by more than 250 eq ha^{-1} yr^{-1}, and these areas are predominantly located in the northeastern US.

293 **Regional patterns in foliar ^{15}N across a gradient of nitrogen deposition in the northeastern US**
L.H. Pardo, S.G. McNulty, J.L. Boggs, S. Duke

Foliar $\delta^{15}N$ increased with increasing N deposition from Maine to NY, but decreased between 1987–1988 and 1999.

303 **Changes in conifer and deciduous forest foliar and forest floor chemistry and basal area tree growth across a nitrogen (N) deposition gradient in the northeastern US**
J.L. Boggs, S.G. McNulty, L.H. Pardo

Data from the 1999 remeasurement of the red spruce forests suggest that N deposition, to some extent, is continuing to influence red spruce across the northeastern US as illustrated by a significant correlation between N deposition and red spruce foliar %N. Our data also suggest that the decrease in forest floor %N and net nitrification potential across sites from 1987 to 1999 may be due to factors other than N deposition, such as climatic factors and N immobilization in fine woody material (>5 cm diameter).

315 **Uncertainty analysis on simple mass balance model to calculate critical loads for soil acidity**
H. Li, S.G. McNulty

A comprehensive uncertainty analysis, with advanced techniques and full list and full value ranges of all individual parameters, was used to examine a simple mass balance model and address questions of error partition and uncertainty reduction in critical acid load estimates that were not fully answered by previous studies.

327 **Influence of ozone pollution and climate variability on net primary productivity and carbon storage in China's grassland ecosystems from 1961 to 2000**
W. Ren, H. Tian, G. Chen, M. Liu, C. Zhang, A.H. Chappelka, S. Pan

Net primary productivity and carbon storage across China's grassland in the late half of the 20th century have been assessed by using the Dynamic Land Ecosystem Model.

336 **Impacts of climatic and atmospheric changes on carbon dynamics in the Great Smoky Mountains National Park**
C. Zhang, H. Tian, A.H. Chappelka, W. Ren, H. Chen, S. Pan, M. Liu, D.M. Styers, G. Chen, Y. Wang

Model simulations suggest that rising atmospheric CO_2 compensates for the adverse effects of ozone stress on ecosystem carbon dynamics in Great Smoky Mountain National Park.

348 **Assessing the risk of foliar injury from ozone on vegetation in parks in the U.S. National Park Service's Vital Signs Network**
R. Kohut

An assessment of the risk of foliar ozone injury on plants was conducted for 269 parks in support of the U.S. National Park Service's Vital Signs Monitoring Network Program.

358 Temporal patterns of foliar ozone symptoms on tall milkweed (*Asclepias exaltata* L.) in Great Smoky Mountains National Park
A.H. Chappelka, G.L. Somers, J.R. Renfro

Ground-level ozone has the potential to cause deleterious effects to tall milkweed growing in Great Smoky Mountains National Park.

366 Isozyme markers associated with O_3 tolerance indicate shift in genetic structure of ponderosa and Jeffrey pine in Sequoia National Park, California
J. Staszak, N.E. Grulke, M.J. Marrett, W. Prus-Glowacki

Genetic variation in isozyme markers associated with ozone tolerance differed between parental trees and their progeny in two closely related species of yellow pine.

I Contents of Volume 149